

On some questions related to the Krichever correspondence

D.V. Osipov, A.B. Zheglov*

Abstract

We investigate various new properties and examples of two-dimensional and one-dimensional Krichever correspondence.

1 Introduction

The aim of this paper is to show several new properties and examples of the two-dimensional Krichever correspondence. On the one hand, these examples make a general picture of the Krichever correspondence more clear. On the other hand, they should show a possible way how to generalize the one-dimensional Krichever correspondence to the case of dimension 2.

Also we give some explicit examples for the one-dimensional Krichever correspondence, which explain some unclearness in the paper [3], and give some new KP-equations which appear as corollaries from generalized KP-hierarchy on two-dimensional local skew fields. These skew fields were first considered in [9] and then classified in [15].

First, we recall that the Krichever map for algebraic curves is the following construction. We start from a data $(C, p, \mathcal{F}, e_p, t)$, where C is an irreducible algebraic curve, p is a smooth k -point on C , t is a local formal parameter at p , $\mathcal{F}$ is a torsion free coherent sheaf of rank r on C , e_p is a local formal trivialization of the sheaf $\mathcal{F}$ at p . We consider the embedding from $H^0(C \setminus p, \mathcal{F})$ into $K_p \otimes \mathcal{F}$. This embedding gives us the map from our data to Fredholm subspaces of the space $k((t))^{\oplus r}$ by identifying $K_p \otimes \mathcal{F} \simeq K_p^{\oplus r}$ under fixed e_p and t . This map is called the Krichever map, see [4], [14], [6].

Now we recall the Krichever map for algebraic surfaces (see [10], [11]). We start from a data $(X, C, p, \mathcal{F}, e_p, t, u)$, where X is a projective irreducible normal algebraic surface, C is an ample Cartier divisor on X , p is a smooth k -point on X and C , $\mathcal{F}$ is a rank r vector bundle on X , e_p is a local formal trivialization of $\mathcal{F}$ at p on X , u and t are local parameters at p such that $u = 0$ is a local equation of C on X near p . By

*supported by DFG-Schwerpunkt "Globale Methoden in der Komplexen Geometrie", by RFBR grant no. 05-01-00455, by grant of Leading Scientific Schools no. 489.2003.1, by INTAS grant 05-100000-8118; besides, the first author was supported by RFBR grant no. 04-01-00702 and by grant of Russian Science Support Foundation.

such data a k -subspace W in the sum of two-dimensional local fields $k((t))((u))^{\oplus r}$ is canonically constructed.

Shortly, this construction consists of the following steps. By the data $p \in C \subset X$ we can canonically build the two-dimensional local field $K_{p,C}$, which is isomorphic to $k((t))((u))$ after fixing local parameters u and t (see [8], [1], [2], [7]). We consider the subfield $K_C \subset K_{p,C}$, which is the completion of the function field on X with respect to the valuation given by the curve C , and the subring $B_p = \varinjlim_n \frac{\hat{\mathcal{O}}_p}{u^n}$, where $\hat{\mathcal{O}}_p$ is the complete local ring at p on X . (B_p does not depend on the choice of u). Now we have

$$W = \bigcap_{q \in C, q \neq p} (B_q \otimes \mathcal{F}) \cap (K_C \otimes \mathcal{F})$$

And from the embedding of the space $K_C \otimes \mathcal{F}$ to the space $K_{p,C} \otimes \mathcal{F}$ we have the embedding of the space W to the space $k((t))((u))^{\oplus r}$ by chosen e_p , t , and u .

Note also that the Krichever map was generalized to algebraic varieties of higher dimensions in the work [5] by higher-dimensional local fields.

We are grateful to A.N. Parshin for the stimulating remarks and discussions. We are grateful also to H. Kurke, and to the Humboldt University of Berlin for the nice atmosphere, where the part of this work was done.

2 Some explicit examples of the two-dimensional Krichever correspondence

In this section we shall compute some examples of the two-dimensional Krichever correspondence on $\mathbf{P}^2$.

In [10] was given an example when $X = \mathbf{P}^2$ with homogeneous coordinates $(x : y : z)$, the curve C is given by $z = 0$, the point p is $(1 : 0 : 0)$, the bundle $\mathcal{F} = \mathcal{O}_X$, the local parameters $t = y/x$, $u = z/x$. Then the subspace $W = k[t^{-1}]((t^{-1}u))$.

For our examples we need the following lemma.

Lemma 1 *Suppose that there is an ample effective divisor D on X such that $D \cdot C = mp$ for some integer m , and the affine curve $C \setminus p$ is given by the equation $f = 0$ on the affine surface $X \setminus D$ for some $f \in k[X \setminus D]$. Choose any $y_1, \dots, y_l \in k[X \setminus D]$ such that their images in $k[C \setminus p]$ generate this ring. (Such y_i always exist). We have that canonically $y_1, \dots, y_l, f \in K_{p,C}$. Then we have*

$$W = k[y_1, \dots, y_l]((f)) \subset K_{p,C}$$

.

Proof follows very easily from the definition of subspace W and facts that $K_C k(C)((f))$ and $B_q = \varinjlim_n \frac{\hat{\mathcal{O}}_q}{f^n}$ for any point $q \in C$, $q \neq p$.

Now we shall apply this lemma to the case $X = \mathbf{P}^2$.

Consider the case of quadric curve.

Proposition 1 Consider the surface $X = \mathbf{P}^2$ with the homogeneous coordinates $(x : y : z)$. Let the curve C be given by the equation $y^2 + z^2 - 2xz = 0$, and the point $p = (1 : 0 : 0)$. Let local parameters $t = y/x$, $u = (y/x)^2 + (z/x)^2 - 2z/x$. Then for the sheaf $\mathcal{F} = \mathcal{O}_X$ we have

$$W = k[\alpha]\left(\left(\frac{\alpha^2}{t^2}u\right)\right),$$

where $\alpha = \frac{t(1+(1+u-t^2)^{1/2})}{t^2-u}$.

Proof. To apply lemma 1, we consider the curve D which is the tangent line to the curve C at the point p . So D is given by the equation $z = 0$, and $D \cdot C = 2p$. Now in the affine domain $z \neq 0$ we have an equation $f = (y/z)^2 + 1 - 2x/z$. Note that $f = ux^2/z^2$, and $k[C \setminus p] = k[y/z]$. Denote $\alpha = y/z$. Then from lemma 1 we obtain that $W = k[\alpha]\left((u\alpha^2/t^2)\right)$. Let us calculate α . From the expression for u we have that $u = t^2 + (t/\alpha)^2 - 2t/\alpha$. Hence $t/\alpha = 1 + (1 + u - t^2)^{1/2}$ or $t/\alpha = 1 - (1 + u - t^2)^{1/2}$. But $t/\alpha = z/x$ has zero of order 2 at the point p after the restriction to the curve C . Therefore for t/α we have the second variant, and hence we obtain the formula from the proposition. The proposition is proved.

Note that for any quadric on $\mathbf{P}^2$ we can take the tangent to the point and apply lemma 1.

Now we consider the case of any curve C of degree m on $\mathbf{P}^2$. Suppose that there exists a point $p \in C$ such that the tangent line D to the curve C at p has the property $D \cdot C = mp$.

For example, if C is a cubic curve, then such point p always exists. (But for the case of a quartic curve this fact is not true anymore, for a general quartic curve such point does not exist.)

Choose homogeneous coordinates $(x : y : z)$ on $\mathbf{P}^2$ such that D is given by $z = 0$, and p is $(1 : 0 : 0)$. Then we have that the affine curve $C \setminus p$ on the affine surface $X \setminus D$ is given by the equation $f(x/z, y/z) = 0$. Therefore we can choose the local parameters $u = (z/x)^m f$ and $t = y/x$. Also the images of elements x/z and y/z generate the ring $k[C \setminus p]$. Hence by lemma 1 for the sheaf $\mathcal{F} = \mathcal{O}_X$ we obtain

$$W = k[\beta, \beta t]\left((\beta^m u)\right), \tag{1}$$

where $\beta = x/z$ is defined from the equation $f(\beta, \beta t) = 0$ and the condition that β has a pole of order m after the restriction to the curve C .

Define the support of a k -subspace W from $k((t))((u))$ as the closed k -subspace of $k((t))((u))$ generated by all monomials which are the lowest degrees of elements from W .

Note that the similar subspaces play important role for the case of Krichever correspondence for algebraic curves. For example, the "holes" between monomials of negative degrees are Weierstrass gaps, their quantity is the genus of the curve. Also in this case it is possible to reconstruct the cohomology groups of original sheaf on the curve from the support of a subspace arising from a geometrical data.

If we are interesting in k -subspaces from $k((t))((u))$ which are images of the Krichever correspondence for algebraic surfaces, it makes sense to investigate the possible supports for such subspaces. But note at once that not all the properties will hold for the case of algebraic surfaces. So, from the support of the subspace we can not reconstruct the cohomology groups of the original sheaf on a surface. (For example, we do not know anything about the cohomology group H^1 from the support of the subspace.)

Now from explicit calculations above we have some examples of subspaces-supports which appear from the Krichever correspondence for algebraic surfaces.

Proposition 2 *Let the conditions be as in formula 1. Then the support of such subspace W is the closed subspace generated by all monomials $t^i u^j$ with $i \leq -m^2 j$ except for the monomials $t^{-\alpha_1 - m^2 l} u^l, \dots, t^{-\alpha_g - m^2 l} u^l$, where i, j and l are any integers, and natural numbers $\alpha_1, \dots, \alpha_g$ are the Weierstrass gaps for the curve from our data.*

Proof follows by the direct application of formula 1. And $\alpha_1, \dots, \alpha_g$ are natural numbers which can not be obtained by linear combinations of integers m and $m - 1$ with integer nonnegative coefficients (g is the genus of the curve).

3 Stabilizer ring of a k -subspace of a two-dimensional local field.

This section is one of the first steps to describe explicitly the images of the two-dimensional Krichever map, that is, to determine k -subspaces of $k((t))((u))$ which are in the image of the two-dimensional Krichever map.

Recall the case of the Krichever correspondence for the curves (see [4]). For any Fredholm k -subspace of $k((t))$, that is, for the k -subspace W such that $\dim_k W \cap \mathcal{O} < \infty$, and $\dim_k k((t))/(W + \mathcal{O}) < \infty$, where $\mathcal{O} = k[[t]]$, we consider the ring

$$A \stackrel{\text{def}}{=} \{a \in k((t)) : a \cdot W \subset W\}.$$

Then the ring A is finitely generated over the field k , and either $A = k$, or the ring A has the Krull dimension equal to 1.

In the second case the subspace W is in the image of the Krichever map, that is, the curve $C = \text{Proj}(\bigoplus_n A \cap t^{-n} \mathcal{O})$, the sheaf $\mathcal{F} = \text{Proj}(\bigoplus_n A \cap t^n \mathcal{O})$, the point p appears from the given filtration $A \cap t^n \mathcal{O}$ of the ring A , t is a formal local parameter at p , and $1 \in k((t))$ is a formal trivialization of $\mathcal{F}$. Moreover, the original k -subspaces A and W appear as the images of the Krichever map of the constructed geometrical data $(C, p, \mathcal{O}_C, 1, t)$ and $(C, p, \mathcal{F}, e_p, t)$ correspondingly. Note that this description is connected with the description of commutative subrings in the ring of differential operators in one variable, that is, with the "Schur pairs", see [4].

Recall that the Euler characteristic of a Fredholm subspace $W \subset k((t))$ is equal to $\chi(W) = \dim_k W \cap \mathcal{O} - \dim_k k((t))/(W + \mathcal{O})$.

Consider now the case of a two-dimensional local field. Recall that a two-dimensional local field has, by definition, two valuations ν and $\bar{\nu}$, where ν is a discrete valuation of

rank 1 on a two-dimensional local field, and $\bar{\nu}$ is a discrete valuation of rank 1 on the residue field of a two-dimensional local field. Let the space $V = k((t))((u))$, and define k -subspaces $\mathcal{O}_1 = k((t))[[u]]$, $\mathcal{O}_2 = k[[t]]((u))$. For any integer n , for any k -subspace $W \subset V$ let $W(n) = (t^n \mathcal{O}_1 \cap W) / (t^{n+1} \mathcal{O}_1 \cap W)$.

For any commutative ring $B \supset k$ we say $\text{trdeg}_k B = m$ if m is the maximal number of algebraically independent elements of the ring B over the field k .

Theorem 1 *Let W be a k -subspace of the space V such that for any integer n the space $W(n)$ is a Fredholm subspace in a one-dimensional local field, and $\chi(W(n)) = a + bn$, where $b < 0$. Let the ring A be a k -subring of the space V such that $A \supset k$, $A \cdot W \subset W$. Then*

- a) for any element $a \in A$ we have $\bar{\nu}(a) \leq b\nu(a)$.*
- b) $\text{trdeg}_k \text{Frac}(A \cap \mathcal{O}_2) \leq 2$, and the field $\text{Frac}(A \cap \mathcal{O}_2)$ is finitely generated over the ground field k . (For any commutative ring without zero divisors by $\text{Frac } B$ we denote its field of fractions.)*

Proof. a) We assume the converse. Then there exists an element $x \in A$ such that $\bar{\nu}(x) > b\nu(x)$. We have $x \cdot W \subset W$ and $x \cdot W(0) \subset W(m)$. It is easy to see that $\chi(x \cdot W(0)) = \chi(W(0)) + \bar{\nu}(x)$. Now we have

$$\chi(W(m)) = a + bm < a + \bar{\nu}(x) = \chi(W(0)) + \bar{\nu}(x) = \chi(x \cdot W(0)) \leq \chi(W(m)).$$

It is a contradiction.

- b) Instead of this case we will prove more general result:

Lemma 2 *Let B be a subring in a two dimensional local field $V = k((t))((u))$ such that $k \subset B$ and the following condition holds:*

for every $a \in B$ we have $0 \leq \bar{\nu}(a) \leq -\pi\nu(a)$, $\pi > 0$.

Then $\text{trdeg}_k \text{Frac } B \leq 2$, and the field $\text{Frac } B$ is finitely generated over the ground field k .

Proof. Consider the subspace $\Delta(N) = \{a \in B | \nu(a) > -N\}$, where the integer $N > 0$. Note that this subspace has a finite dimension over the field k , and this dimension is not greater than πN^2 .

Indeed, if this dimension is greater than πN^2 , then the ring B must contain an element x such that $\bar{\nu}(x) > -\pi\nu(x)$. It is a contradiction.

Note that $\text{trdeg}_k B < 2$ if and only if $\bar{\nu}(a) = l\nu(a)$ for some constant l and all elements $a \in B$. Indeed, if there are two elements a, b such that the last condition does not hold, then they must be algebraically independent, because the first monomials of such elements are algebraically independent. Converse, if $\bar{\nu}(a) = l\nu(a)$ for all $a \in B$, then the supports (the monomials of lowest degrees) of all elements from B lie on a line which goes through the zero point of a coordinate plane, and therefore one can apply arguments, for example, from [4, Prop. 3.2.] to obtain $\text{trdeg}_k B < 2$.

We suppose now that $\text{trdeg}_k B \geq 2$. We take two elements $a, b \in B$ such that $\bar{\nu}(a)/\nu(a) \neq \bar{\nu}(b)/\nu(b)$. It means that their supports do not belong to a line which goes

through the zero point on a coordinate plane. These elements exist because $\text{trdeg}_k B \geq 2$, and because of the arguments from the previous paragraph. Also, they are algebraically independent elements. We can estimate the dimension of a subspace generated over the field k by powers of elements a, b which lie in the subspace $\Delta(N)$ for some N . It is clear that this dimension is greater than

$$\frac{1}{2} \cdot \left(-\frac{N}{\nu(a)} - 1\right) \left(-\frac{N}{\nu(b)} - 1\right) = eN^2 + lN + r$$

for some real numbers e, l, r and for every sufficiently large number N . (To obtain this estimate we consider a parallelogram constructed by vectors which go from the zero point of a coordinate plane and end in the points $a^{[-N/\nu(a)]}$ and $b^{[-N/\nu(b)]}$ correspondingly.)

Now suppose that the field $\text{Frac } B$ has infinite dimension over the field $\text{Frac } k[a, b]$. Then we take M linearly independent over the ring $k[a, b]$ elements $a_1, \dots, a_M$, where the number M satisfies the condition $Me > \pi$. Without loss of generality we can assume that $\nu(a_1) \leq \nu(a_i)$ for all i . Since the dimension of a subspace generated over the field k by powers of elements a, b which lie in the subspace $\Delta(N + M\nu(a_1))$ is greater than $e(N + M\nu(a_1))^2 + l(N + M\nu(a_1)) + r$ for every sufficiently large number N , we obtain that the dimension of a subspace generated over the field k by elements $a_1, \dots, a_M$ multiplied by powers of elements a, b which lie in the subspace $\Delta(N + M\nu(a_1))$ is greater than $M(e(N + M\nu(a_1))^2 + l(N + M\nu(a_1)) + r)$ for every sufficiently large number N . The last subspace is inside the space $\Delta(N)$. On the other hand, since $Me > \pi$, for every sufficiently large number N we have

$$M(e(N + M\nu(a_1))^2 + l(N + M\nu(a_1)) + r) > \pi N^2 > \dim_k \Delta(N).$$

We obtained a contradiction. Therefore, in this case the field $\text{Frac } B$ has finite dimension over the field $\text{Frac } k[a, b]$. Hence, $\text{trdeg}_k \text{Frac } B = 2$, and the field $\text{Frac } B$ is finitely generated over the ground field k .

The lemma is proved.

The theorem is proved.

Remark 1. As we have seen in the proof of the lemma, $\text{trdeg}_k B = 1$ if and only if the supports of all elements from the ring B lie on a line which goes through zero point on a coordinate plane. (Moreover, arguing similar to [4, Prop. 3.2.]), it is possible to prove that the ring B is finitely generated over the field k and has Krull dimension equal 1.)

If $\text{trdeg}_k B = 0$, then $B = k$.

Remark 2. It is important that in theorem 1 we consider the intersection $A \cap \mathcal{O}_2$. The statement b) of theorem 1 is not true for the ring A itself as the following example shows.

Consider the subspace $W = \{a \in V \mid \bar{\nu}(a) \leq -\nu(a)\}$. It is possible to check that the subspace W satisfies the conditions of theorem 1, and that $A = W$. But since the ring A contains the subfield $k((ut^{-1}))$, we have that the field $\text{Frac } A$ has the infinite transcendency degree over the field k .

Remark 3. In theorem 1 we consider the ring $A \cap \mathcal{O}_2$, because this ring is the image of the affine ring $H^0(X \setminus C, \mathcal{O}_X)$ when the pair (A, W) is the image of the geometrical data $(X, C, p, \mathcal{O}_X, 1, t, u)$, $(X, C, p, \mathcal{F}, e_p, t, u)$ under the Krichever map. And we need exactly this ring for the theorem of reconstruction of geometrical data. Namely, the required surface $X = Proj(\oplus_n (A \cap \mathcal{O}_2) \cap t^n \mathcal{O}_2)$, the required curve C appears from the filtration induced by the filtration $t^n \mathcal{O}_2$ on $A \cap \mathcal{O}_2$, and the required point p appears from the one-dimensional Krichever correspondence on the curve C , which is given by $W(0)$ (see [10]).

Also the space $W(n)$ is the image of the sheaf $\mathcal{F}|_C \otimes N_{C/X}^{\otimes -n}$ under the Krichever map on the curve C . And by the Riemann-Roch theorem applied to the curve C we obtain the Euler characteristic

$$\chi(N_{C/X}^{\otimes -n}) = 1 - g(C) + \deg(\mathcal{F}) - n \deg N_{C/X}.$$

Besides $N_{C/X} = \mathcal{O}(C \cdot C)$. And from ampleness of the divisor C we have $\deg N_{C/X} > 0$. It explains the condition $\chi(W(n)) = a + bn$, $b < 0$ from theorem 1.

Remark 4. If a k -subspace W is the image of a geometrical data $(X, C, p, \mathcal{F}, e_p, t, u)$ under the Krichever map, then from the complex which calculates the cohomology groups (see [10], [11]) we have

$$\begin{aligned} W \cap \mathcal{O}_1 \cap \mathcal{O}_2 &= H^0(X, \mathcal{F}) \\ \frac{W \cap (\mathcal{O}_1 + \mathcal{O}_2)}{W \cap \mathcal{O}_1 + W \cap \mathcal{O}_2} &= H^1(X, \mathcal{F}) \\ \frac{V}{W + \mathcal{O}_1 + \mathcal{O}_2} &= H^2(X, \mathcal{F}). \end{aligned}$$

And we know that the dimensions of these subspaces over the field k are finite as dimensions of cohomology groups. It would be interesting to add these conditions of finite-dimensionality to conditions of theorem 1 and to obtain new corollaries for the answer, if a k -subspace $W \subset V$ is in the image of geometrical data under the Krichever map?

4 To the question about an embedding of a projective line into an infinite-dimensional affine space

In the paper [3, p. 13] was remarked the following curious fact: there is an embedding of the universal Sato Grassmannian into an infinite-dimensional affine space. In this section we clarify, if this embedding is algebraic. To this end we investigate an example, in which this embedding are restricted to the projective line in the universal Sato Grassmannian.

First we recall some well-known facts about the Sato Grassmannian, and also the construction of an embedding map from the paper [3].

Let $V = k((z))$ be the field of Laurent power series with a filtration $V(n) = z^n k[[z]]$. Let $V_1 = V(0)$.

By $Gr(V)$ we denote the set of subspaces W in V such that the complex

$$W \oplus V_1 \rightarrow V$$

is a Fredholm one. This set possess the structure of an infinite-dimensional projective variety, whose connected components are marked by the value of the Euler characteristic of the complex. We will work only with "zero" component $Gr_0(V)$.

Now we consider the skew field $P = k((x))((\partial^{-1}))$ of formal pseudo-differential operators with coefficients from the field $k((x))$. This skew field is a left $k((x))$ -module of all expressions $L = \sum_{i>\infty}^n a_i \partial^i$, $a_i \in k((x))$ with a multiplication defined according to the Leibnitz rule.

In the same way one can define the ring $E = k[[x]]((\partial^{-1}))$, $E \subset P$. It can be checked that P and E are associative rings (see details in the paper [9]).

There is a decomposition

$$E = E_+ + E_-,$$

where $E_- = \{L \in E : L = \sum_{n<0} a_n \partial^n\}$, and E_+ consists of the operators containing only nonnegative powers of ∂ . The elements from $E_+ =: D$ are the usual differential operators, and the elements from E_- are the Volterra operators.

The map $E \rightarrow E/Ex = V$ (we identify the image of ∂^{-1} with z) defines a linear action of the ring E on V and therefore on $Gr(V)$. The map $E \rightarrow V$ is called *the Sato map*.

Let us introduce the notion of a standard subspace:

Definition 1 Let S be a subset in $\mathbb{Z}$, $S := \{\sigma(0), \sigma(-1), \sigma(-2), \dots\}$, where $\sigma(-i) \in \mathbb{Z}$, $\sigma(-i) = -i$ for $i \gg 0$.

The subspaces $V^S := \bigoplus_{l \in \mathbb{N} \cup 0} k \cdot z^{\sigma(-l)} \subset V$, $V^S \in Gr_0(V)$ are called the standard subspaces.

By W_0 we denote the subspace V^{S_0} , where $S_0 = \{-1, -2, \dots\}$.

The following lemma was proved in the paper [13].

Lemma 3 There is a unique operator R such that $RV^S = W_0$. Namely,

$$R = \partial^{-m-1}(x\partial - \sigma(0))(x\partial - (\sigma(-1) + 1))(x\partial - (\sigma(-2) + 2)) \dots (x\partial - (\sigma(-m) + m)),$$

where m is the maximal number such that $\sigma(-m) - m \neq 0$.

The notion of a quasiregular operator was introduced in the paper [12].

Definition 2 Let $\mathcal{E}_K^{(0)}$, $K = k((x))$ be a group of monic (i.e., with the leading coefficient equal 1) operators of degree zero from the ring P .

An operator $W \in \mathcal{E}_K^{(0)}$ is called quasiregular if there are numbers $m, n \in \mathbb{N}$ such that $x^m W$ and $W^{-1}x^n$ belong to E .

By $\mathcal{R}$ we denote the set of all quasiregular operators.

Now we can introduce a map $\gamma : \mathcal{R} \rightarrow Gr_0(V)$. Put

$$\gamma(W) = (W^{-1}x^n)W_0, \quad W \in \mathcal{R}.$$

Since $x \cdot W_0 = W_0$, this definition is correct, i.e., it does not depend on n . It is clear that $\gamma(W) \in Gr_0(V)$.

The following theorem was proved in the paper [13].

Theorem 2 *The map $\gamma : \mathcal{R} \rightarrow Gr_0(V)$ is a bijection.*

It was remarked in the paper [3, p. 13] that $\mathcal{E}_K^{(0)}$ is an infinite dimensional affine space, and $\mathcal{R}$ is embedded into this space. (It is more correctly to say that $\mathcal{E}_K^{(0)}$ is an inductive limit of infinite-dimensional affine spaces). It was also remarked that due to the theorem there is an embedding of the universal Sato Grassmannian into this affine subspace. To check this assertion we give below an explicit computation of this embedding for a projective line which lies in the Sato Grassmannian.

Consider the set of Fredholm subspaces

$$R = \{W(\alpha, \beta) \subset V, W(\alpha, \beta) = \bigoplus_{l=1}^{\infty} k \cdot z^{-l} \oplus k \cdot (\alpha + \beta z)\},$$

where $\alpha, \beta \in k$. It is clear that R is a projective line in $Gr_0(V)$ with coordinates $(\alpha : \beta)$.

Consider the operators $\tilde{S}(\alpha, \beta) = \alpha + \beta x + \beta \partial^{-1}$. We have $\tilde{S}(\alpha, \beta) \in E$. Note that $\tilde{S}(\alpha, \beta)W_0 = W(\alpha, \beta)$ for any pair (α, β) . In the case $\alpha \neq 0$ we define the operator $S(\alpha, \beta) = \alpha^{-1}\tilde{S}(\alpha, \beta)(1 + \beta/\alpha x)^{-1}$. Since $(1 + \beta/\alpha x)W_0 = W_0$, we have $S(\alpha, \beta)W_0 = W(\alpha, \beta)$. Note that $S \in \mathcal{E}_K^{(0)} \cap E$. In the case $\alpha = 0$ it is possible to represent the operator $\beta^{-1}\tilde{S}(\alpha, \beta)$ in the form $H(\alpha, \beta)^{-1}x$, where $H(\alpha, \beta) \in \mathcal{E}_K^{(0)}$, $H(\alpha, \beta) = (1 + \partial^{-1}x^{-1})^{-1} = 1 - x^{-1}\partial^{-1} + \dots$. Thus, we obtain

$$\gamma^{-1}(W(\alpha, \beta)) = \begin{cases} S(\alpha, \beta)^{-1} = 1 - (\frac{\beta}{\alpha} - \frac{\beta}{\alpha}x + \frac{\beta^2}{\alpha^2}x^2 + \dots)\partial^{-1} + \dots & \text{if } \alpha \neq 0 \\ H(\alpha, \beta) = 1 - x^{-1}\partial^{-1} + \dots & \text{if } \alpha = 0. \end{cases}$$

Now consider two coordinate functions: the affine space $\mathcal{E}_K^{(0)}$ has coordinates a_{ij} , where a_{ij} are coefficients of the series $1 + \sum_{i,j} a_{ij}u^i\partial^{-j}$ (any element from $\mathcal{E}_K^{(0)}$). For our projective line we have the following coordinates under the embedding:

$$a_{-1,1} = \begin{cases} 0 & \text{if } \alpha \neq 0 \\ -1 & \text{if } \alpha = 0 \end{cases}$$

$$a_{0,1} = \begin{cases} -\beta/\alpha & \text{if } \alpha \neq 0 \\ 0 & \text{if } \alpha = 0. \end{cases}$$

These functions are not continuous! Hence, it follows that the embedding of the universal Sato Grassmannian into the affine space is only a set-theoretic map.

5 New equations of KP-type on skew fields

In this section we give the answer on a question set up in the paper [9]. We recall that the classical KP-hierarchy is constructed by means of the ring of pseudo-differential operators $P = k((x))((\partial^{-1}))$. This ring is a skew field. The point is to consider other skew fields instead of this one. We study below, if there are some new non-trivial generalizations of the KP-hierarchy for a known list from the classification of two-dimensional local skew fields. In particular, we give a number of new equations of the KP-type.

We recall that in the paper [9] A.N.Parshin pointed out a class of non-commutative local fields and showed that these skew fields possess many properties of commutative fields. He defined a skew field of formal pseudo-differential operators in n variables and studied some of their properties. He raised a problem of the classification of all non-commutative local skew fields. In the paper [15] this problem was solved for $n = 2$. A.B. Zheglov obtained the list of skew fields up to an isomorphism. Among other skew fields this list contains a classical ring of pseudo-differential operators P mentioned in the previous part of this paper. The following theorem was proved in [15]:

Theorem 3 (I) *Let K be a two-dimensional local skew field with a commutative residue skew field.*

It splits¹ if the canonical automorphism² α satisfies the condition $\alpha^n \neq Id$ for all n . If this condition does not hold, there are examples of non-splittable skew fields.

(II) *Let K, K' be skew fields as in (I). Assume that $\alpha^n \neq Id$ for all n . Then*

(a) *K is isomorphic to a two-dimensional local skew field $\bar{K}((z))$, where $za = a^\alpha z$, $a \in \bar{K}$ and $\bar{K}$ is a one-dimensional local field with the residue field k .*

(b) *K and K' are isomorphic if and only if $k \cong k'$, and there is an isomorphism $f : \bar{K} \mapsto \bar{K}'$ such that $\alpha = f^{-1}\alpha'f$.*

(c) *If $\text{char} K = \text{char} k$, $\text{char} K' = \text{char} k'$ and k, k' are algebraically closed fields of characteristic 0, then K is isomorphic to K' if and only if $k \cong k'$ and $(a_1, i_\alpha, y(\alpha)) = (a'_1, i_{\alpha'}, y(\alpha'))$.*

(III) *Let K, K' be splittable two-dimensional local skew fields of characteristic 0, $k \subset Z(K)$, $k' \subset Z(K')$, and $\alpha^n = Id$, $\alpha'^n = Id$ for some natural numbers $n, n' \geq 1$. Then*

(a) *K is isomorphic to a two-dimensional local skew field $k((u))((z))$, where*

$$zuz^{-1} = \xi u + u^{\delta'_{i_n}} z^{i_n} + u^{\delta'_{2i_n}} z^{2i_n},$$

where $\xi^n = 1$, $i_n = i_n(0, \dots, 0)$,

$\delta'_{i_n}(u) = cu^{r_n}$, $c \in k^/(k^*)^e$, $e = (r_n - 1, i_n)$,*

$\delta'_{2i_n}(u) = (a_n(0, \dots, 0) + r_n(i_n + 1)/2)u^{-1}(\delta'_{i_n}(u))^2$

(i_n, r_n, a_n were defined in [15]).

If $n = 1$, $i_n = \infty$, then K is commutative.

(b) *K is isomorphic to K' if and only if $k \cong k'$ and the sets $(n, \xi, i_n, r_n, c, a_n)$, $(n', \xi', i'_n, r'_n, c', a'_n)$ coincide.*

Corollary 1 *Every two-dimensional local skew field K with the set*

$$(n, \xi, i_n, r_n, c, a_n)$$

¹a local skew field is called splittable if there is a section of the residue homomorphism

²the canonical automorphism of a local skew field is defined as an automorphism of the residue skew field induced by the inner automorphism $Ad(z)$ of the skew field, where z is an arbitrary parameter

is a finite-dimensional extension of the skew field with the set $(1, 1, 1, 0, 1, a)$.

For every skew field from this list it is possible to define a decomposition $K = K_+ + K_-$, where $K_- = \{L \in K : \text{ord}(L) < 0\}$ ³, and K_+ consists of the operators containing only nonpositive (≤ 0) powers of z , and a "KP-hierarchy" in the Lax form:

$$\frac{\partial L}{\partial t_n} = [(L^n)_+, L],$$

where $L \in z^{-1} + K_- \otimes k[[\dots, t_m, \dots]]$. Let $L = z^{-1} + u_1 z + u_2 z^2 + \dots$, where $u_m = u_m(u, t_1, t_2, \dots)$. In the sequel we will write $\partial/\partial t_n$ as ∂_n .

One can easily check that if the canonical automorphism α from the classification theorem 3 is not trivial, then our "KP-hierarchy" becomes trivial in a sense that it can be easily linearized and solvable. We omit calculations here. Thus, we can assume that $\alpha = \text{id}$. In this case, if $i > 1$, then we have $[(L^n)_+, L] = -[(L^n)_-, L] = 0 \pmod{\wp^i}$, where $\wp$ is the ideal of the first valuation in K . Then, as before, our "KP-hierarchy" is linear and easily solvable.

So, we have to consider the case $i = 1$. In this case we have $r = 0$ and $c = 1$ (see [15]), and, therefore, there is only one non-trivial parameter a . If $a = 0$, then K is isomorphic to the ring P of usual pseudo-differential operators under $x = -u$, $\partial = z^{-1}$. By $u'_i, u''_i, \dots$ we denote the subsequent derivatives by x .

For $n = 1$ we obtain the following equation

$$\partial_1 u_1 = u'_1,$$

i.e., we can assume that $t_1 = x$ for u_1 .

Now we write down the first two equations for $n = 2$ and the first equation for $n = 3$.

$$\partial_2 u_1 = u''_1 + 2u'_2 \quad (2)$$

$$\partial_2 u_2 = 2u'_3 + 2u_1 u'_1 + u''_2 - a(2x^{-1}u'_2 + 2x^{-1}u''_1 - x^{-2}u'_1) \quad (3)$$

$$\partial_3 u_1 = u'''_1 + 3u''_2 + 3u'_3 + 6u_1 u'_1 - 3a(x^{-1}u''_1 - x^{-2}u'_1). \quad (4)$$

We introduce the following notation: $u = u_1(x, y, t)$, where $y = t_2$, $t = t_3$. We will use also the standard notations $u_t, u_y, u_{yy}, \dots$ for partial derivatives.

Eliminate u'_3 from equations (3) and (4). We obtain

$$3u_{2y} - 2u_t = -6uu' - 3u''_2 - 2u''' - 3a(2x^{-1}u'_2 + x^{-2}u'). \quad (5)$$

From equation (2) we find

$$u'''_2 = 1/2(u''_y - u''''_y), \quad u'_{2y} = 1/2(u_{yy} - u''_{yy}).$$

Differentiating equation (5) by x and using these expressions we finally obtain a new nontrivial KP-equation:

$$(4u_t - u''' - 12uu')' = 3u_{yy} + 6a(2x^{-2}u'' - x^{-2}u_y - x^{-1}u''' + x^{-1}u'_y - 2x^{-3}u').$$

³we use $\text{ord}(\sum_k a_k z^k) = -\min\{k : a_k \neq 0\}$

It is easily to see that if $a = 0$, then we have the usual KP-equation (compare also explicit calculations in the paper [11]).

Other interesting generalizations of KP-hierarhies which are connected with the development of Krichever correspondence to higher dimensions (sections 1-3) are described in [16].

References

- [1] Fimmel T., Parshin A.N. *An introduction to the higher adelic theory*, preprint.
- [2] Huber A., *On the Parshin-Beilinson Adeles for Schemes*, Abh. Math. Sem. Univ. Hamburg, **61** (1991), 249-273.
- [3] Mulase M., *Solvability of the super KP equation and a generalization of the Birkhoff decomposition*, Inv. Math., Vol. 92, Fasc. 1, 1988, 1-47.
- [4] Mulase M., *Category of vector bundles on algebraic curves and infinite dimensional Grassmannians*, Int. J. Math., **1** (1990), 293-342.
- [5] Osipov D. V., *Krichever correspondence for algebraic varieties*, Izv. Ross. Akad. Nauk Ser. Mat. 65 (2001), no. 5, 91–128; english translation in Izv. Math. 65 (2001), no. 5, 941–975; see also e-print math.AG/0003188.
- [6] Osipov D. V., *The infinite-dimensional Sato Grassmannian and coherent sheaves of rank 2 on curves*, Mat. Sb. 194 (2003), no. 11, 81–94; english translation in Sb. Math. 194 (2003), no. 11-12, 1665–1678; see also e-print math.AG/9904152.
- [7] Osipov D. V., *n-dimensional local fields and adeles on n-dimensional schemes*, to appear in Lecture Notes Series of London Mathematical Society; see also e-print math.AG/0508205
- [8] Parshin A. N. *On the arithmetic of two-dimensional schemes, I. Repartitions and residues*, Izv. Akad. Nauk SSSR Ser. Mat. **40** (1976), no. 4, 736-773; English transl. in Math. USSR Izv. **10** (1976), no. 4, 695-729.
- [9] Parshin A. N., *On a ring of formal pseudo-differential operators*, Proc. Steklov Math. Institute, **224** (1999), 266-280, see also e-print math.AG/9911098
- [10] Parshin A. N., *Krichever correspondence for algebraic surfaces*, Funct. Anal. Appl., v.35 (2001), n.1, pp. 88-90, see also extended version in e-print alg-geom/9911097.
- [11] Parshin A.N., *Integrable systems and local fields*, Communications in Algebra, 29(9), 4157-4181 (2001).
- [12] Sato M., Sato Y., *Soliton Equations as Dynamical Systems on Infinite Dimensional Grassmann Manifold*, Lecture Notes in Num. Appl. Anal., **5** (1982), 259-271.

- [13] Sato M. and Noumi M., *Soliton equations and universal Grassmann manifold* (in Japanese), Sophia Univ. Lec. Notes Ser. in Math. 18 (1984), pp. 74-75
- [14] Segal G., Wilson G. *Loop groups and equations of the KdV type*, Publ. Math. IHES, 80(1985), 301-342.
- [15] Zheglov A.B. *On the structure of two-dimensional local skew fields*, Izvestiya RAN: Ser. Mat., 1, 2001, 25-60 in russian; English transl. in Izvestiya: Mathematics **65:1** (2001) 23-55.
- [16] Zheglov A.B. *Two dimensional KP systems and their solvability*, e-print math-ph/0503067

D.V. Osipov
 Steklov Mathematical Institute RAS
 e-mail *d_osipov@mi.ras.ru*

A.B. Zheglov
 Moscow State Lomonosov University
 e-mail *azheglov@mathematik.hu – berlin.de*